

# Pleomorphic Adenoma of Buccal Mucosa: An Uncommon Presentation among Cheek Swellings

**Sailesh KM<sup>1</sup>, Amit K<sup>2</sup>, Ejaz Ahmad M<sup>2</sup> and Shweta P<sup>2</sup>**

Department of Dentistry, AIIMS, India

Resident Doctor, Department of Dentistry, AIIMS, India

**\*Corresponding author:** Sailesh KM, Assistant Professor, HOD Department of Dentistry, AIIMS, Bihar, India, E-mail: dr.saileshmukul@yahoo.co.in

## Case Report

Volume 2 Issue 2

**Received Date:** April 21, 2017

**Published Date:** May 13, 2017

**DOI:** 10.23880/oajds-16000134

## Abstract

Cheek mass is a common presenting problem in Oral & Maxillofacial OPD. It may be due to many underlying causes like inflammation, cyst or neoplasm. Common cheek swellings sometimes come out as an uncommon lesion on histopathological examination where the provisional diagnosis is challenged. Therefore the final diagnosis of common small cheek swelling is not complete till its therapeutic removal and histopathological examination. We represent a case of young healthy female who reported with swelling of right side buccal mucosa which was finally diagnosed as pleomorphic adenoma based on histopathologic examination.

**Keywords:** Pleomorphic Adenoma; Minor salivary gland tumor; Cheek mass

## Introduction

Among all head and neck neoplasm, Salivary gland neoplasm represents 3-5% [1] and minor salivary glands tumor account for about 22% of all salivary gland neoplasm [2]. Majority of minor salivary gland tumors reported in literature are malignant with only 18% being benign [3]. Pleomorphic adenoma (PA) is a benign, mixed tumor which most commonly involves the parotid gland. Among tumors of the minor salivary glands palate is the most common intra-oral site (42.8-68.8%) followed by upper lip (10.1%) and cheek (5.5%). Other rare sites include the throat (2.5%), retro molar region (0.7%), alveolar mucosa and floor of the mouth [4]. Vaidya, et al. [5] conducted a study of 104 minor salivary gland tumor and concluded that there was no gender predilection and the median age of presentation was 45 years [5]. Minor salivary gland tumors are rare in children. They are responsible for only 5-10% of all salivary tumors under 20 years of age [6-8]. The presentation of Pleomorphic adenoma is as a painless, well-delineated, slow growing smooth, firm, mobile swelling that does not cause

ulceration of overlying mucosa [9]. This paper describes the diagnosis and management of an asymptomatic slowly growing pleomorphic adenoma of buccal mucosa in a young patient. It is suggested that a practitioner has to be aware of such a rare condition.

## Case Report

A 19 year old female patient reported to the department with chief complains of unaesthetically looking painless swelling of right side cheek since one year. The swelling started as a painless mass which gradually increased to the presenting size of 2cm X 1.5 cm. the size remained constant in last 4 months. The swelling was present over the right cheek region at infero anterior border of the mandible. The skin over the anteroinferior aspect of the swelling had blind spot (punctum) (Figure 1). On palpation swelling was soft, non-compressible, non-tender, and non-pulsatile and attached to overlying skin. Swelling was not appreciable

intra orally. No signs of inflammation were present. Ultrasound (USG) of the swelling revealed well defined cystic lesion with echogenic content and multiple small anechoic areas, no vascularity was present and provisional diagnosis of sebaceous cyst was made. Surgical excision of lesion was planned under GA. Through intra oral approach an incision was made over the mucosa, the underlying muscle was dissected and the mass was approached. The mass was dissected free from overlying skin. The mass was encapsulated with small multilobulated surface and pale yellowish appearance (Figures 2 & 3). Histopathology report showed areas of chondromyxoid stroma, proliferating ductal epithelial cells, myoepithelial cells, squamous differentiation with keratinization. Histological features were consistent with Pleomorphic Adenoma.



Figure 1: The swelling presented as a painless mass of size of 2 cm X 1.5 c. The skin over the anteroinferior aspect of the swelling had blind spot (punctum).



Figure 2: The mass was encapsulated with small multilobulated surface and pale yellowish appearance.



Figure 3: The mass was dissected free from overlying skin.

## Discussion

WHO pathological classification of salivary gland tumors has proved to be most useful in daily practice and has been adopted worldwide [10]. Pleomorphic adenomas are benign heterogeneous tumors of salivary gland origin. In head and neck region they arise mainly in the major salivary glands mostly parotid (85%), submandibular gland (8%) and intra oral (7%) salivary glands. They are commonly seen in 4 -6 decades of life [8]. The etiology of pleomorphic adenoma is not known. It is of epithelial origin and cytogenetical analyses of pleomorphic adenomas have shown the presence of clonal chromosomal aberrations on the basis of which

## Classification in Three Cytogenetic Subgroups have been Made

- I. Pleomorphic adenomas with structural chromosomal aberrations of chromosomes 8 in the region 8q12.
- II. Ii) Pleomorphic adenoma with structural aberrations of chromosome 12 in the region 12q 13-15.
- III. Iii) Pleomorphic adenomas with normal karyotypes possessing also the cell lines with the trisomy 8 or the loss of chromosome Y as mosaicism [11].

Histopathological features of pleomorphic adenoma are a variable pattern of epithelium in a loosely fibrous stroma which may be myxoid, chondroid or mucoid. The epithelium is usually arranged in sheets or strands, and ductal structures, often bilayered, are typical. In the minor glands, lesions are often more solid or cellular than those seen in the major glands, and the myoepithelial cells are often polygonal with a pale eosinophilic cytoplasm giving plasmacytoid an epithelioid phenotype [12] as present in

our case. The main histologic subgroups have been identified: myxoid (80% stroma), cellular (myoepithelial predominant) and mixed (classic) type [13]. Rarely, a malignant tumor may arise within this tumor, a phenomena known as carcinoma ex Pleomorphic Adenoma [14]. PA should be considered in the differential diagnosis of mass of the cheek in youngsters [15]. Differential diagnosis of intraoral, solid, asymptomatic swelling include dermoid cyst, neurofibroma, rhabdomyosarcoma, lipoma, foreign body reaction, buccal space abscess, mucocele, fibroma, adenoid cystic carcinoma and mucoepidermoid carcinoma [6,16]. The ideal treatment for PA is wide local excision with good safety margins and regular follow up for 3-4 years [17]. Since these tumors are radio resistant, the use of radiation therapy is contraindicated.

To conclude we can say that Pleomorphic adenoma of buccal mucosa is an uncommon provisional diagnosis. It must be considered in differential diagnosis for cheek mass in youngsters as its complete resection with surrounding dispensable normal tissue is paramount. Recurrence after long period of excision and malignant transformation is a concern and therefore long period of follow up is necessary.

## References

1. Leegaard T, Lindeman H (1969) Salivary-gland tumours clinical picture and treatment. *Acta Otolaryngol Suppl* 263: 155-159.
2. Thoeny HC (2007) Imaging of salivary gland tumours. *Cancer Imaging* 7(1): 52-62.
3. Hakeem AH, Hazarika B, Pradhan SA (2009) Primary pleomorphic adenoma of minor salivary gland in the parapharyngeal space. *World J Surg Oncol* 7: 85.
4. Ca W, Sk M, Gnepp DR (1988) Tumors of the intraoral minor salivary glands: a demographic and histologic study of 426 cases. *Oral Surg Oral Med Oral Pathol* 66(3): 323-333.
5. Vaidya AD, Pantvaidya GH, Metgudmath R, Kane SV, D'Cruz AK (2012) Minor salivary gland tumors of the oral cavity: a case series with review of literature. *J Cancer Res Ther* 1:111-115.
6. Lotufo MA, Junior CAL, Mattos JP De, França CM (2008) Pleomorphic adenoma of the upper lip in a child. *J oral Sci* 50(2): 225-228.
7. Jorge J, Pires FR, Alves FA, Perez DEC, Kowalski LP, et al. (2002) Juvenile intraoral pleomorphic adenoma: report of five cases and review of the literature. *Int J Oral Maxillofac Surg* 31(3): 273-275.
8. Dalati T, Hussein MR (2009) Juvenile pleomorphic adenoma of the cheek: a case report and review of literature. *Diagn Pathol* 4: 32.
9. Toida M, Shimokawa K, Makita H, Kato K, Kobayashi A, et al. (2005) Intraoral minor salivary gland tumors: a clinicopathological study of 82 cases. *Int J Oral Maxillofac Surg* 34(5): 528-532.
10. Yousef V, Mohammad Ali M (1995) Review of 103 Cases of Minor Salivary Gland Tumors. *Acta Medica Iranica* 33(1-2): 22-26.
11. Bullerdiel J, Bartnitzke S (1991) Chromosomal abnormalities in pleomorphic adenomas: Related to origin or karyotypic evolution? *Cancer Genet Cytogenet* 53(2): 283-284.
12. Speight PM (2007) Update on Diagnostic Difficulties in Lesions of the Minor Salivary Glands. *Head Neck Pathol* 1(1): 55-60.
13. Ogata H, Ebihara S, Mukai K (1994) Salivary gland neoplasms in children. *Jpn J Clin Oncol* 24(2): 88-93.
14. Goyal P, Sehgal S, Ghosh S, Agrawal D, Singh S (2016) Rare carcinoma ex-pleomorphic adenoma of buccal mucosa: case report and review of literature. *Rare Tumors* 8(1): 6138.
15. Mk Gupta, Bailoor DN, Mhaske S, Raghuvanshi V, Nahar S, et al. (2013) Juvenile Pleomorphic Adenoma of the Cheek - A rare case with review of literature. *J Dent Med Sci* 6(1): 45-48.
16. Yamamoto H, Fukumoto M, Yamaguchi F, Sakata K, Oikawa T (1986) Pleomorphic adenoma of the buccal gland in a child. *Int J Oral Maxillofac Surg* 15(4): 474-477.
17. Amit A, Ravinder S, Soheyl S, Shambulingappa P, Isha S (2012) Pleomorphic adenoma of minor salivary gland: a case report. 9(1): 97-101.

